

Authors' affiliations

F Boccara, A Cohen, Department of Cardiology, Saint Antoine University Hospital, Assistance Publique-Hôpitaux de Paris and Université Paris VI, Paris, France

E Teiger, Department of Physiopathology, Henri Mondor University Hospital, Assistance Publique-Hôpitaux de Paris and Université Paris XII, Creteil, France

REFERENCES

- Haffner SM, Lehto S, Ronnemaa T, *et al*. Mortality from coronary heart disease in subjects with type 2 diabetes and in nondiabetic subjects with and without prior myocardial infarction. *N Engl J Med* 1998;**339**:229–34.
- Gu K, Cowie CC, Harris MI. Diabetes and decline in heart disease mortality in US adults. *JAMA* 1999;**281**:1291–7.
- Berger PB, Velianou JL, Aslanidou Vlachos H, *et al* for the BARI Investigators. Survival following coronary angioplasty versus coronary artery bypass surgery in anatomic subsets in which coronary artery bypass surgery improves survival compared with medical therapy. Results from the bypass angioplasty revascularization investigation (BARI). *J Am Coll Cardiol* 2001;**38**:1440–9.
- Legrand V, Serruys PW MD, Unger F, *et al* on behalf of the Arterial Revascularization Therapy Study (ARTS) Investigators. Three-year outcome after coronary stenting versus bypass surgery for the treatment of multivessel disease. *Circulation* 2004;**109**:1114–20.
- Holmes DR Jr, Vlietstra RE, Smith HC, *et al*. Restenosis after percutaneous transluminal coronary angioplasty (PTCA): a report from the PTCA Registry of the National Heart, Lung, and Blood Institute. *Am J Cardiol* 1984;**53**:77C–81C.
- Van Belle E, Bauters C, Hubert E, *et al*. Restenosis rates in diabetic patients: a comparison of coronary stenting and balloon angioplasty in native coronary vessels. *Circulation* 1997;**96**:1454–60.
- Van Belle E, Perie M, Braune D, *et al*. Effects of coronary stenting on vessel patency and long-term clinical outcome after percutaneous coronary revascularization in diabetic patients. *J Am Coll Cardiol* 2002;**40**:410–7.
- Sabate M, Jimenez-Quevedo P, Angiolillo DJ, *et al* for the DIABETES Investigators. Randomized comparison of sirolimus-eluting stent versus standard stent for percutaneous coronary revascularization in diabetic patients: the diabetes and sirolimus-eluting stent (DIABETES) trial. *Circulation* 2005;**112**:2175–83.
- Roiron C, Sanchez P, Bouzamondo A, *et al*. Drug eluting stents: an updated meta-analysis of randomised controlled trials. *Heart* 2006;**92**:641–9.
- Kastrati A, Dibra A, Eberle S, *et al*. Sirolimus-eluting stents vs paclitaxel-eluting stents in patients with coronary artery disease: meta-analysis of randomized trials. *JAMA* 2005;**294**:819–25.
- Stettler C, Allemann S, Egger M, *et al*. Efficacy of drug eluting stents in patients with and without diabetes mellitus: indirect comparison of controlled trials. *Heart* 2006;**92**:650–7.
- Dibra A, Kastrati A, Mehilli J, *et al* ISAR-DIABETES Study Investigators. Paclitaxel-eluting or sirolimus-eluting stents to prevent restenosis in diabetic patients. *N Engl J Med* 2005;**353**:663–70.
- Morice M-C, Serruys PW, Colombo A, *et al*. Eight-month outcome of the REALITY study: a prospective, randomized, multicenter head-to-head comparison of the sirolimus-eluting stent (Cypher) and the paclitaxel-eluting stent (Taxus). Presented at the 2005 Annual Scientific Session of the American College of Cardiology, Orlando, Florida, March 6–9, 2005.
- Windecker S, Remondino A, Eberli FR, *et al*. Sirolimus-eluting and paclitaxel-eluting stents for coronary revascularization. *N Engl J Med* 2005;**353**:653–62.
- Moliterno DJ. Healing Achilles—sirolimus versus paclitaxel. *N Engl J Med* 2005;**353**:724–7.
- Iakovou I, Schmidt T, Bonizzoni E, *et al*. Incidence, predictors, and outcome of thrombosis after successful implantation of drug-eluting stents. *JAMA* 2005;**293**:2126–30.
- Colombo A, Cosgrave J. Paclitaxel-eluting stents in complex lesions. *JAMA* 2005;**294**:1268–70.

IMAGES IN CARDIOLOGY

doi: 10.1136/hrt.2005.071365

Double coarctation and double grafts: role of computed tomographic angiography in diagnosis of a clinic mystery

A 27 year old woman was admitted to our clinic with left upper extremity pain of two months' duration. History of the patient revealed two operations for coarctation of the aorta. The left subclavian artery to descending thoracic aorta and the ascending aorta to descending thoracic aorta grafts were interposed at 2002 and 2003, respectively. Blood pressure was 120/80 mm Hg from the right brachial artery, 70/40 mm Hg from the left brachial artery, and 90/60 mm Hg from the left and right popliteal arteries. Auscultation revealed second degree systolic ejection murmur at the left upper sternal border radiating to the interscapular area. To define the underlying cause of the extremity pain computed tomographic (CT) angiography was performed. Three dimensional CT angiography (see panel) identified the proximal (asterisk) and distal (double asterisk) coarctation segments, the two grafts, and the occluded left subclavian artery. Unfortunately, the first graft (arrowhead) was interposed between the left subclavian artery and the descending aorta in between the two coarctation segments which, possibly, made a second operation necessary. The second graft (arrow) was interposed between the ascending aorta (Asc. Ao) and descending thoracic aorta (Desc. Ao). Shunting of blood via the aorta–aortic graft and the recent operation may be the underlying causes of the left subclavian artery occlusion and left upper extremity pain. CT angiography provides suitable images for the diagnosis and localisation of coarctation of the aorta, which may aid in surgical planning.



A Yildiz
S Ozturk
G Cagirci
ghcayildiz@yahoo.com